

The Anthropocene Shield: Melanin-Driven Resilience against CFC-Induced Ozone Volatility

Marie-Soleil Seshat Landry, CEO of Landry Industries
ORCID: 0009-0008-5027-3337

January 9, 2026

Abstract

This paper models the survival advantage of high-eumelanin phenotypes in the context of the Ozone Depletion Era. We analyze the catalytic destruction of O_3 by Chlorofluorocarbons (CFCs) and the subsequent biological impact on Folate (B_9) stability.

1 The Chemical Catalyst of Vulnerability

The release of Freon (CF_2Cl_2) into the stratosphere facilitates the catalytic breakdown of ozone. The resulting increase in UVB flux (Φ_{UVB}) is inversely proportional to the total column ozone (Ω):

$$\Phi_{UVB} \propto \frac{1}{\Omega} \quad (1)$$

2 The Biological Shield (Eumelanin)

Melanin provides a protective coefficient ($\mathcal{P}$) that prevents the photolysis of 5-MTHF. The preservation of Folate $[F]$ over time t is modeled as:

$$[F]_t = [F]_0 \cdot e^{-k \cdot \Phi_{UVB}(1-\mathcal{M})} \quad (2)$$

Where $\mathcal{M}$ represents the melanin density. For *TypeVI* skin, $\mathcal{M} \rightarrow 1$, ensuring systemic stability in high-radiation environments.

3 Conclusion

The Anthropocene favors the melanin-rich. Strategic biological sovereignty requires recognizing this protection as a primary asset.