

PUBLIC SERVICE ANNOUNCEMENT: The "Ringer" Is a Warning

Author: Marie-Soleil Seshat Landry

Date: March 4, 2026

Harm Reduction | Acute Hypertension | Vasoconstriction | Tinnitus | Cardiovascular Risk | Stimulant Toxicity

AI Disclosure: Draft supported by AI language model. Medical claims aligned with peer-reviewed toxicology and public health literature.

1. The Signal

In stimulant intoxication, sudden ear ringing ("the ringer") is not a bonus effect. It is consistent with acute physiological stress. Cocaine sharply increases blood pressure, heart rate, and vascular constriction. The inner ear relies on delicate end-arterial blood supply. Reduced flow can trigger transient tinnitus or auditory distortion.

CLINICAL REALITY: There is no medically established "safe dose" of cocaine. Stroke, seizure, and cardiac arrest have

occurred in small or first-time exposures.

2. What Is Likely Happening Biologically

Cocaine blocks reuptake of dopamine, norepinephrine, and serotonin, producing intense sympathetic activation. This may result in:

- Sudden elevation in blood pressure
- Coronary and cerebral vasospasm
- Increased clotting tendency
- Reduced seizure threshold
- Transient cochlear ischemia

The auditory system is highly sensitive to vascular instability. Ringing can reflect temporary disruption of blood flow or neural hyperexcitability.

3. Risk Minimization Principles

Abstinence eliminates risk. When abstinence is not the immediate choice, public health guidance focuses on reducing catastrophic outcomes — not optimizing intoxication.

Evidence-Based Harm Reduction Includes:

- Avoid mixing with alcohol (increases cardiotoxicity through cocaethylene formation)
- Avoid mixing with opioids (fentanyl contamination risk is significant)
- Do not use alone
- Access drug checking services where available
- Maintain hydration and recovery sleep

- Keep naloxone accessible due to contamination risk

4. Red Flag Symptoms — Seek Immediate Medical Care

- Chest pain or pressure
- Severe or sudden headache
- Weakness on one side of the body
- Confusion or difficulty speaking
- Irregular heartbeat
- Persistent ringing with dizziness

These may signal stroke, myocardial infarction, hypertensive crisis, or acute toxicity.

5. Strategic Perspective

The nervous system has warning mechanisms. Ignoring them does not build resilience; it compounds risk. Chasing threshold experiences is a human pattern. Biology, however, has fixed limits.

The intelligent objective is preservation of cognitive and cardiovascular integrity.

References & Authoritative Sources

1. National Institute on Drug Abuse (NIDA) – Cocaine Research Reports
2. American Heart Association – Cocaine and Cardiovascular Complications

3. Centers for Disease Control and Prevention – Stimulant Overdose Data
4. World Health Organization – Harm Reduction Guidance
5. New England Journal of Medicine – Cocaine-Induced Myocardial Infarction
6. British Medical Journal – Acute Stimulant Toxicity Management
7. Journal of Hypertension – Acute Blood Pressure Effects of Cocaine
8. American Journal of Otolaryngology – Tinnitus and Substance Exposure
9. UNODC – World Drug Report
10. International Journal of Drug Policy – Evidence-Based Harm Reduction

© 2026 Landry Industries Intelligence For educational and public health purposes only. If you or someone you know is struggling with substance use, contact local health services or a confidential support line.